
EVALUATION OF THE PHYSICO-CHEMICAL PROPERTIES OF WATER IN THE CHURU TEHSIL RAJASTHAN (INDIA)

DR. RAKESH KUMAR

Assistant Professor

J J T University Jhunjhunu, Rajasthan

ABSTRACT

Water is an essential natural resource for sustaining life and environment but over the last few decades the water quality is deteriorating due to its over exploitation. Water quality is essential parameter to be studied when the overall focus is sustainable development keeping mankind at focal point. Groundwater is the major source of drinking water in rural as well as in urban areas and over 94% of the drinking water demand is met by groundwater. Ground water is the main source of drinking water in the Churu. Drinking water contamination by fluoride, chloride, nitrate, arsenic etc. play important role in public health. The main health risks from fluoride are bone and teeth diseases. Nitrate can undergo endogenous reduction to nitrite and nitrosation of nitrites can form N-nitroso compounds which are potent carcinogens. Nitrites can lead among infants to the disease called blue baby syndrome. There is strong relation between nitrate concentration and diarrhea and also to some other diseases.

KEYWORDS: Alkalinity, Hardness.

INTRODUCTION

Nelson et al. (2004), says that WQI is defined as an evaluation of the physical, chemical and biological nature of water with respect to its natural quality, human effects and intended uses. It reduces a list of parameters to a simpler expression to enable easier interpretation of monitoring data. Insaf et al. (2007), worked on groundwater quality index using GIS, where he calculated the normalised sub indices and then ranked them. His study area is a few km² and the parameters he used are only chemical parameters containing ions. The final index is represented in % not in values. He then selected the combination of parameters to show the variability of groundwater.

The provision of safe supply of drinking water is a high priority issue to a government for safeguarding the health of human beings. The production and supply of adequate and safe drinking water is the most important factor contributing to a decrease in morbidity and mortality in developing country like India. As we know water is a very good solvent, hence some toxic and hazardous substances dissolve in it and produce water pollution problem posing many parameters of interest for water quality assessment. In Rajasthan 71% of the irrigation and 90% of the drinking water supply sources are underground water (Rathore, 2005).

Churu district is the part of great Indian Thar desert and lies in arid zone of western Rajasthan. The region is situated between 73° 5' to 75°5' E longitude and 27°5' to 29° 0' N latitude. It is bordered by Sri Ganganagar in the north; Nagaur, Sikar and Jhunjhunu in south; Bikaner in the west and Haryana in the east. The physiography of whole district is characterized by shifting and stable sand dunes, which give it a desolate and barren look. Being situated in a shallow depression, the climatic conditions are very extreme here and very hot summer and very cool winter prevail. The rainfall in the district is scanty and irregular. Annual rainfall is very low, which range around 225 to 500 mm (Singh *et al.* 2009).

In absence of surface water resources, people of the area primarily depend upon underground water to meet out their drinking and agricultural requirements. The underground water in the study area is characterized by medium to high salinity. More than 57% water is of sodium chloride type and distributed throughout the district. The district is notorious for having high fluoride and nitrate concentration in underground water, which makes it unsuitable for drinking purpose. The dry climatic condition with high evaporation and

insignificant recharge might have accelerated the strengthening of fluoride and nitrate concentrations in the underground water of this area.

There are many factors contributing to nitrate contamination to water sources in the region. An increase of nitrates in water is often associated with use of agricultural chemical by farmers such as fertilizer, pesticides or poor sanitary activities. It was found that the crop of family legumineaceae is the main reason of nitrate hazard in the Churu. The crop of this family takes atmospheric nitrogen and fixed it in the soil as nitrate and more nitrates are formed than the plants need. Such excess nitrates are available for leaching and ground water get contaminated with high concentration of nitrate. Nitrate is the ultimate oxidation state of nitrogen and is the source of pollution in ground water. Nitrates undergo reduction and nitrosation of nitrite can form N- nitroso compound that is a potent carcinogen. The use of High nitrate contaminated drinking water is a well known risk factor for infant methemoglobinemia (Mirvish, 1985). Gastric cancer, birth defects and hypertrophy of the thyroid are other health effects on humans that are potentially influenced by elevated levels of nitrate in drinking water (Kugali *et al.*, 2013). The increased nitrate level in drinking water may also adversely affect the central nervous system.

The health problems arising as a result of fluoride contamination is more widespread. According to a report of national geophysical Research institute of Hyderabad about 177 districts of India are fluoride affected. The country has increasing incidences of fluorosis, both dental and Skeletal with some 66 million people being at risk (Sinha and Mustaria 2004). In India, Rajasthan has maximum area affected by excess concentration of fluoride in ground water (Murlidharan *et al.* 2002). About 22 districts of Rajasthan have greater amount of fluoride than the permissible limits (Sinha *et al.* 1970). The reason of origin of these hazardous elements is attributed to geological reason. According to recommendation of WHO drinking water should not contain more than 1.5 mg/l fluoride. People who drink water containing fluoride in excess of this level over many years could get bone disease, including pain and tenderness of the bones. In moderate forms of dental fluorosis, brown staining or pitting of permanent teeth may appear.

Alkalinity is the measure of capacity of water to neutralize the acids. It provides guidance in applying doses of chemicals in water and waste water processes specially in coagulations, softening and operational control of anaerobic digestion.

Hardness in ground water in the area is due to availability of limestone, sedimentary rock and calcium bearing minerals in soil. Hardness can also occur locally in groundwater from excessive application of lime to the soil in agricultural areas. Excessive hardness in drinking water results in urinary concentrations, stomach disorder, stone in kidney and bladder.

Chlorides are widely distributed in nature as salts of sodium chloride, potassium chloride, and calcium chloride. A normal adult human body contains approximately 81.7 g chloride. Excessive intake of drinking-water containing sodium chloride at concentrations above 2.5 g/l has been reported to produce hypertension (Fadeeva, 1971). This effect is believed to be related to the sodium ion concentration.

REVIEW OF LITERATURE

Chaudhary *et al.* (2009) assessed fluoride toxicity level in Gang canal catchment area of north-western Rajasthan. Fluoride concentration was found to vary from 0.50 to 8.50 mg/l. in Sujangarg Tehsil of Churu district, ground water analysis revealed 56.94% of the samples had fluoride beyond permissible limit and highest level recoded was 45.02 mg/l (Jangid, 2008). Epidemiological study revealed that 56.37% respondents have severed skeletal fluorosis.

Singh p *et al.* (2011) analyzed the concentration of fluoride in ground water of Churu city. Majority of ground water sample of this area were found to contain much higher fluoride content, beyond the permissible

limits. Maximum concentration (8.5 ppm) occurred at Madersa, (7.8 ppm) at Van Vihar and (7.3 ppm) at Eye's Hospital's Tube well. As a result the leeching out of fluoride from the above source contaminates water and soil. During the study it was also found the water from the deep well had more fluoride contamination as compare to relatively less deep wells. Trivedy et al. (1984) have noted that well waters in all the three towns of Satara district of Maharashtra to be rich in solids, carbonates, total hardness, calcium and magnesium in comparison to the surface waters. Lakshmanan et al. (1985) have reported that concentration of uranium to be marginally higher in well water (5.8 mg/l) than bore well waters (4.2 mg/l). Trivedy and Mathur (1986) have noted very high nitrate content ranging from 5.0 mg/l to 1400 mg/l in ground waters of arid region of Churu district in Rajasthan. Trivedy and Goel (1986) have reported that the hardness is the property of water which prevents the lather formation with soap and increases the boiling points of water.

Govindan and Thanasekaran (1993) have suggested routine monitoring and periodical evaluation as a measure for control of ground and surface water quality in urban and rural water supply scheme. Pandey et al. (1993) studied the sanitary water quality of Bhimtal lake of Kumaun hill. They reported that sodium, potassium and calcium were maximum during Sept. 1990 and decreased steadily during water. Dissolved oxygen, total alkalinity, total solid and total dissolved solids were inversely proportional to BOD and COD. Nag and Das (1995) have studied the drinking water quality in Puruliya district of west Bengal and reported sodium, calcium, chromium, copper, lead, mercury and arsenic below their permissible level but the presence of potassium, magnesium, manganese, iron, zinc and cadmium above their maximum admissible concentration or guideline value of WHO in several samples.

Nag and Das (1994) have studied the drinking water quality in the Birbhum district of West Bengal. They reported that Na, Ca, Mg, Cr, Pb, Hg, and as are below their permissible limits and K, Cu and Cd are above their permissible limits. Altman and Parizek (1995) have reported that there are various ways as ground water is contaminated such as use of fertilizer in farming.

Nawlakhe et al. (1995) have reported that the ground water quality of rural areas in Shivpuri district of Madhya Pradesh and observed that nitrate fluoride, TDS and hardness to be the problem parameters in order of dominance. They suggested reverse Osmosis, electro dialysis, in exchange and solar distillation to be effective process for removal of such multiple problem parameters.

MATERIAL AND METHODS

Total 21 groundwater samples were collected from tube wells, open wells, and hand pumps of different locations of study area. The collected samples were analyzed for estimation of physical and chemical properties like pH, Total Alkalinity, total hardness, Ca, TDS, Chloride, Nitrate and Fluoride. Samples were collected in clean bottles of 1 liter capacity. Physical parameters like, pH, TDS were measured using digital meters immediately after sampling. The total hardness, alkalinity, nitrate, nitrite, chloride, fluoride were analyzed using standard titrimetric methods - APHA 1998.

RESULTS AND DISCUSSION

Present study is an overview of drinking water parameters in Churu and surrounding area. The most of the water samples in study area are brackish to saline in nature and pH values ranged from 7.1 to 8.4, which is more than the WHO (1984) permissible limits. Alkalinity and pH in the groundwater of study area are caused by bicarbonate and carbonates. The values of alkalinity in all the samples range from 180 to 700 mg/l in the area.

The hardness of ground water ranged from 200 to 1480 mg CaCO_3/l with an average of 761.42 mg CaCO_3/l in the study area, whereas WHO permissible limit for hardness of ground water for safe life is 500mg/l. The analysis reveals the increase in the concentration is mainly due to mineralization of ground water. The range

of calcium in the groundwater of study area varied from 120 to 700 mg/l. The amount of TDS in groundwater depends on the solubility of minerals in different geological regions of the area. Water containing TDS concentrations below 1000 mg/l is usually acceptable for consumption. In the present study TDS was found to vary from 920 to 4620 mg/l.

Chloride concentration range was from 200 to 2550 mg/l in the study area, which is higher in the most of samples than desirable level determined by ICMR (200 mg/l). Except three, all the study sites had chloride contents far above the permissible limit of WHO. Chloride concentration was maximum in Balrasar and minimum in Gajsar.

High concentration of nitrate has been reported in ground water of Churu. Data analysis reveals that nitrate concentration in collected water samples lie in the range from 30 to 280 mg/l, with an average of 81.19 mg/l (Table 1). WHO proposed permissible limit for nitrate is 45 mg/l but about 19% of samples collected in study area have high values of nitrate. Churu (GPF) and Balrasar had maximum nitrate contents far above the WHO permissible limit.

Fluoride content ranges from 0.5 to 4.3 mg/l in the Churu tehsil with an average of 1.66 mg/l (Table 1). The study indicates that most of the water samples contain high concentration of fluoride than that of the standard set by WHO and ICMR (1.5 mg/l). Churu T/W 31 had the maximum fluoride contamination followed by Churu GPF site. Ten study sites had fluoride contamination above WHO permissible limit and rest had the concentration within range. The analysis of data indicates that high concentration of fluoride is associated with salinity. Regular intake of fluoride contaminated water may cause bone diseases.

The results of data analysis showed that there is a significant association between chloride, fluoride and nitrate contaminated drinking water and diseases related to bone, teeth and digestion

Table-1
Physico-chemical parameters of groundwater in selected areas of Churu Tehsil

S. No.	Location (Churu Tehsil)	pH	Alkalinity	Hardness		TDS	Chloride	Nitrate	Fluoride
				Total	Calcium				
1	Asloo	7.6	300	1200	700	4620	2000	30	1.9
2	Balrasar	7.6	420	1420	720	2700	2520	230	1.7
3	Churu (GPF)	7.5	240	410	170	920	250	280	3.1
4	Churu Garh	7.8	430	1070	120	2960	800	50	0.9
5	Churu Goenka School	7.4	400	630	250	1725	600	30	1.0
6	Churu Mishrawell	7.5	630	360	200	2480	700	45	2.6
7	Churu T/W-3	7.7	620	580	170	2960	900	50	1.9
8	Churu T/W-31	7.7	640	460	130	2350	750	40	4.3
9	Churu T/W-58	7.8	700	670	230	2900	1050	50	2.4
10	Churu T/W-95	7.6	500	690	290	2760	750	50	1.8
11	Dhadhar T/W-I	7.7	680	640	320	2680	1380	125	1.5
12	Dhadhar T/W-II	7.1	680	580	310	3050	1580	120	1.6
13	Dhadhria charnan	7.5	380	960	460	3090	1700	90	0.5
14	Gajsar road	7.8	300	200	110	1020	200	30	1.3
15	Ghanghu	7.1	240	1480	620	1940	620	95	1.4
16	Khansoli	8.0	450	930	450	3720	1580	55	1.2
17	Lakhau	7.7	280	1000	570	4340	1780	65	1.7
18	Pithisar	8.4	180	1260	680	1940	580	70	0.9
19	Raipuria	7.5	360	580	280	2940	900	50	1.4
20	Ratan nagar	7.8	440	310	180	1760	610	65	1.1
21	Shyopura	7.4	480	560	370	2770	780	80	0.7

The quality of groundwater of Churu and surrounding area is not according to norms of Rajasthan government and WHO standards. Earlier studies conducted in the Sadulpur tehsil of the district also revealed more or less same results (Singh 2014). Higher concentration of these salts had deleterious effect on the human health (Fetter 1980). Higher concentration of nitrates had been found to be associated with cancer in Slovakia (Gulis *et al.* 2002). Knobetocho (2000) reported blue baby syndrome associated with high nitrate concentration in water. Neal *et al.* (2000) analyzed the water quality of Thames tributaries and found them to be contaminated by anthropogenic activities. Arsenic contamination has found to be lethal in many areas of India (Chakraborti *et al.* 2004, Rahman *et al.* 2005) and Nepal (Shretha *et al.* 2003). Fortunately arsenic contamination is not very high in the region. Akhilesh *et al.* (2005) carried out study on the geochemical scenario of fluoride in the whole Rajasthan. The pH results indicate that ground water of Churu tehsil is highly brackish to saline in nature. The total hardness of ground water of this area falls in the hard category. Salinity, hardness, chloride, nitrates and fluoride properties of drinking water in the study area indicate the sign of deterioration which calls for at least primary treatment of ground water before being used for drinking. During sample collection people of area also complained about diarrhea, gastric and other digestion related problems, which may be because of consuming ground water containing high amount of nitrate contents. Consumption of high concentration of fluoride has resulted in bone deformity and joint pains. Laanthe *et al.* (2002) has suggested some bioremediation methods of nitrate but these have yet to gain fruitful results.

The purpose to persuade present research work was to know ground water quality and to find out chemical and biochemical remediation of problematic ions. The study suggests that regular evaluation of water quality is needed in Churu tehsil so that strategies can be developed to reduce excess chemical contaminations and the water can be made suitable for drinking purposes.

REFERENCES

1. Arif, M. Hussain, I. Hussain, J. Kumar, S. 2012. Fluoride Distribution in Ground Water and Survey of Dental Fluorosis in Villages of Didwana Tehsil of Nagaur District of Central Rajasthan, India. India Water Week 2012- Water, Energy and Food Security.
2. Hussain I, Arif M, Hussain J. (2011), Fluoride Contamination in Drinking water in Rural Habitations of Central Rajasthan, India, Environmental Monitoring and Assessment (DOI: 10.1007/s10661-011-2329-7) Online Published on 20 September, 2011)
3. Keshar Dev (2014). Estimation of the Physico-Chemical Properties of Water in the Churu Tehsil, of Arid Region of Rajasthan (India), Vol.-III, Issue-III, July-2014, RNI No. UPENG/2012/42622
4. Malik, S., A. Singh and V. Menon (2009). Assessment of water quality of upper lake, Bhopal (M.P.). Proc. National Sem. on Sci. Edu. and Attraction of Talent for excellence in Res., held on 22-22 feb., at Bhopal (M.P.). pp. 73.
5. McLay, C.D.A., R. Dragten, G. Sparling, N. Selvarazah (2001). Predicting ground water nitrate concentration in a region of mixed agricultural land use: A comparison of three approaches. Environ Pollut. 115: 191- 204.
6. Mitharwal S., Yadav R.D., and Angasaria R.C. (2009), Water Quality analysis in Pilani of Jhunjhunu District (Rajasthan)- The place of Birla's Origin. Rasayan Journal of Chemistry. 2(4), pp 920-923.
7. Mukul Bishnoi and S. Arara (2007). Potable ground water quality in some village of Haryana, India: Focus on fluoride. J. Environ. Biol., 28: 291-294.
8. Robertson, W.D., J.A. Cherry E.A. Sudicky (1991). Ground water contamination from two small septic systems in sand aquifers, J. Ground water. 29(1): 82-91.
9. Saxena, S., Saxena, U. (2013), Study of fluoride contamination status of ground water in Bassi Tehsil of district Jaipur, Rajasthan, India. International Journal of Environmental Sciences.3(6): 2251-2260.
10. Sharma, N., A.Joram, S. Sharma (2009). Study of water quality parameters in Makarera fresh water lake, near Bewar (Ajmer). Proc. of National Conference on Environ. Health Hazards Held on 17th to 18th Dec. at Kota (Raj.). ID : 7895, PP. 60-61.
11. Smitha, P.G., K. Byrappa and S.N. Ramaswamy (2007). Physicochemical characteristics of water sample of Bantwal Taluk, S. Western Karnataka, India. J. Environ. Biol., 28(3): 591-595.
12. Susana, J.D.P., J. Paggi and P. Collins (2008). Water quality and zooplankton composition in receiving pond of the storm water run off from an urban catchment. J. Environ. Bio., 29 (5).
13. Venkatasubramani, R. and T. Meenambal(2007). Study of surface water quality in Mattupalayam Taluk of Coimbatore District, Tamilnadu, Nat. Environ. Pollut. Tech., 6: 307-310.