

COMPARATIVE EXAMINATIONS OF THE BIOCHEMICAL COMPONENTS OF SEVERAL CULTIVARS OF MUNG BEANS GROWN IN RAINFED AREAS

Irfanul Hasson

Research Scholar, Glocal School of Science
The Glocal University, Mirzapure Pole , Saharanpur (U.P).

Prof. (Dr.) Satyavir Singh

Research Supervisor, Glocal School of Science
The Glocal University Mirzapure Pole , Saharanpur(U.P).

Abstract

The field experiment to determine Biochemical, Molecular and Morphological Characterization of Mung bean [*Vigna radiata* (L.) Wilczek] Cultivars in Rainfed Area” of Uttar Pradesh” during kharif and jayad season in the year 2021-22 to 2022-23 was conducted at the Glocal University, Saharanpur, Uttar Pradesh. The nutritional composition of *Vigna radiata* were estimated in terms of carbohydrate, free amino acid, protein, crude protein, methionine, fat content and nutrient content such as Fe, Cu and Zn biochemical changes in seeds. The significantly maximum (63.56%) carbohydrate in the variety of RMG-62 but maximum total soluble sugar content (6.11%) and minimum nutrient value noticed in Pant M-2. These results suggest that the nutritional content and quality of *Vigna radiata* significantly differ according to cultivars.

Introduction

In India, pulses find an important place in human diet due to vegetarian food habits. According to FAO estimate, 70 per cent of human food comprises cereals and legumes and the remaining 30 per cent comes from animal. The area under pulses in India is around 24.38 million hectares with a production of 14.52 million tones (Sathyamoorthiet al., 2017). Nearly 8 per cent of this area is occupied by mungbean, which is the third most important pulse crop of India, next to gram and pigeonpea. In Punjab, mungbean occupied 12.3 thousand hectares with a production of 10.7 thousand tonnes and with average productivity of 873 kg per hectare (Anonymous, 2018). Pulses are generally deficient in sulfur containing amino acids such as methionine and cysteine but rich in lysine and tryptophan (Khan et al., 2019), which are the limiting amino acids in cereals. Mungbean seeds contain 22-25% protein, 1.21% lysine, 0.006% iron and other essential elements. The seeds are used as whole as well as split dal, which is easy to cook. Nutrients, hormones and environmental factors such as temperature, relative humidity may alter biochemical components affecting seed quality and quantity. A lot of work on growth parameters such as short stature, short duration, synchronous maturity, shining green bold seeds, resistance to shattering of pods, high grain yield and tolerance to Mungbean Yellow Mosaic Virus (MYMV) has been reported (Sekhon et al., 2004). Mungbean is considered the hardiest of all pulse crops since

it is grown extensively under varying climatic conditions. It has greater capacity to grow at high temperature up to 45°C. Therefore, presence of hard seeds affects the grain quality and yield. Work on mungbean biochemical components, storage protein, their characterization, electrophoretically and immunologically for subunit composition and changes during seed development has been reported (Sital et al., 2009). Protein quality improvement in mungbean as influenced by sulfur nutrition and Rhizobium inoculation has also been studied. Schroeder(2016) compared some characters pertaining to protein quality parameters viz; crude protein, extractable protein, globulins, albumins, total seed and protein sulfur, carbon: nitrogen and nitrogen: sulfur ratios in the genus *Pisum*, but information in comparison to climatic conditions affecting seed biochemical components of mungbean was lacking. Therefore, the present investigation was carried out to comparative studies on biochemical components in mung bean [*Vigna radiata* (L.) Wilczek] varieties cultivars in rainfed area during Kharif and Jayad seasons in Bhopal region.

Materials and Methods

The current study was carried out at an agricultural farm of the School of Agricultural science, Glocal University University during the Kharif of 2021–22 and 2022–23 U.P. (India). The area is characterised by a dry sub-humid climate. A set of twelve cultivars of mungbean [*Vigna radiata* (L.) Wilczek] (BPMR-145, HUM-2, MUM-2, PAIRY MUNG, PANT M-2, PANT M-4, PKV AKM-4, PUSA-0672, RMG-62, RMG-268 and SAMRAT were taken as experimental materials in the present study. Three replications of each cultivar were used in the experiment's Randomized Block Design (RBD) layout during the Kharif seasons of 2020–21 and 2021–22. The field tests took place between July and September of 2020–21 and 2021–22. Each genotype was planted on an area of 3.0 m by 2.25 m and having five rows. The plant-to-plant spacing was kept at 10 cm by thinning, and the row-to-row spacing was 45 cm. A sample of mungbean seeds weighing 0.1 g was extracted twice using 5 ml of hot, 80% ethanol. The reducing sugar was dissolved in 10ml of distilled water after the supernatant was collected and evaporated on a water bath at 80°C. Then, after adding 3 ml of the dinitrosalicylic acid (DNS) reagent to a 1 ml aliquot, the mixture was heated on a water bath for 5 minutes. 1 ml of 40% Sodium - Potassium tartarate was added while the tube's contents were still heated and then cooled to room temperature. At 510 nm, a spectrophotometer (Systronics 169) was used to measure the intensity of the red colour in comparison to a reagent blank. On the basis of a standard curve created with known concentrations of glucose, the total amount of reducing sugar was estimated. The estimation was carried out three times, and the mean of the results was expressed as a percentage of reducing sugar in the moisture-free sample. The data were statistically analyzed using factorial complete random design and mean values of different parameters were compared using critical difference at 5%.

Results and Discussion

The largest figure for moisture percent was found in HUM-2 (10.67%) in the pooled data, along with Pant M-2 (9.43%) and RMG-268 (9.23%), and the least value was discovered in Samrat (7.83%). The outcomes were consistent with those of Blessing and Gregory (2010) and Mubarak (2005), who estimated that raw, unprocessed mungbean flour had a moisture content of 10.25% and 9.75%, respectively. The moisture content of mungbean flour is 8.78% in Afzal (1978) and 8.25% in Bhatta et al. (2000), respectively. The variations in moisture content may be influenced by geographical and varietal factors. The majority of nutritional calories

consumed by humans and other animals come from carbohydrates. The combined data also showed that MUM-2 (62.47%) had the highest carbohydrate content, followed by Samrat (62.09 %) and Pusa-9072 (62.05%), while HUM-2 (59.67%) had the lowest value. The findings of this study corroborate those of Mubarak (2005), Agugo and Onimawo (2008), Paul et al. (2011), who determined that the carbohydrate content was 62.9%, 61.47%, and 60.35 percent, respectively. According to Adel et al. (1980), the percentage of carbohydrate in mungbean seeds ranged from 64.15 to 66.32%. Mungbean seeds have a total carbohydrate content of 54.9–58.9% of their weight, according to Habbibullah et al. (2007). Savage and Deo and Muller (1988) and Savage and Deo and Muller (2000) both noted significantly reduced carbohydrate content (between 35 and 40 percent). Different chemical synthesis due to environmental and genetic variables may be the source of the variation in total carbohydrate content. The combined data also showed that Samrat (4.47%) had the highest total soluble sugar content, while RMG-62 (3.97 %) had the lowest. HUM-2 (3.97%) and Pant M-4 (2.19%) were the next-highest in this list. In contrast to the most recent findings, Naivikul and D'apponia (1976) showed that sugar content was higher in mungbean genotypes and was 7.22 percent higher. According to Anonymous (2012), soluble sugar concentrations differed among mungbean genotypes (between 7.1% and 8.9%). According to Mondal et al. (2013), the soluble sugar concentration of freshly harvested seeds from six different mungbean kinds ranged from 5.01 to 9.30%. The combined data also showed that BPMR-145 (3.96%) had the highest non-reducing sugar value, followed by Pusa-9072 (3.17%) and Samrat (2.98%), while Pant M-2 (2.17%) had the lowest value. In both the years. Non-reducing sugar in mungbean, broadbean, and kidneybean varied from 5% to 7%, according to Tanusi et al. in 1972. On the other hand, a substantially lower range, 7.10–7.11 mg/100g mungbean flour, has been observed by (Kakati et al. 2010). The combined results also showed that the cultivar HUM-2 had the highest reducing sugar content (0.92%), followed by the cultivars Pant M-2 (0.80%) and Samrat (0.80%), while cultivar Pusa-0672 had the lowest (0.39%). These findings are in line with those of Chakraborty (1993) and Kakati et al. (2010), who discovered that the decreasing sugar levels in mungbean flour varied from 641.61 to 794.50 mg/100g and from 724.97 to 729.23 mg/100g, respectively, on a dry weight basis. Tanusi et al. measured decreasing sugar levels in mungbean, broadbean, and kidneybean in 1972; these levels ranged from 0.06 to 0.10 g per 100g on a dry weight basis. Mubarak (2005) found an unusually high result (4.85 g/100g) for reducing sugar in raw mungbean seed during his investigation of the effects of different domestic and traditional methods on the nutritional composition of mungbean seeds. The combined data also showed that HUM-2 (25.16%), BPMR-145 (23.98%), and Pusa-0672 (24.97%) had the highest and lowest values of crude protein, Soluble Protein Content, Methionine Content, Crude Fat Content, Calorific Value, Total Phenol Content, Ash Content respectively. Samrat (23.00%) had the lowest value. In 1998, Saleem et al. showed that the crude protein content of mungbean seed ranged from 22.88 to 24.65 percent. The percentages of protein in mungbean seed flour reported by Agugo and Onimawo (2008), Blessing and Gregory (2010), Butt and Batool (2010), and Gregory and Gregory (2008) are 25.09%, 24.08%, 25.90%, and 25.00%, respectively. (2000) Bhatti et al. The variations in protein levels are assumed to be caused by the genetic makeup of mungbean cultivars, cultural practices, as well as some environmental factors.

The cultivar BPMR-145 (14.89 mg/100g) had the highest quantity of iron, followed by Pusa-9072 (14.28 mg/100g) and PKV AKM-4 (14.12 mg/100g), while Pusa-0672 (10.32 mg/100g) had the lowest amount. The combined results showed that Pusa-9072 had the highest copper and zinc concentration, followed by RMG-62, and HUM-2, while Samrat had the lowest value (1.55 mg/100g). Habibullah et al. (2007) found that two kinds

of mungbean had zinc concentrations that were much lower (1.54-1.88 mg/100 g flour).

Table: 1 Values of Moisture (%), Total Carbohydrate (%), Total Soluble Sugar (%), Non Reducing Sugar(%) in 12 Mung bean Cultivars

| S. N. | Cultivar/Variety | Moisture (%) | Total Carbohydrate (%) | Total Soluble Sugar (%) | Non Reducing Sugar (%) |
|-------|-------------------|--------------|------------------------|-------------------------|------------------------|
| 1 | BPMR-145 | 8.2 | 61.54 | 2.99 | 3.96 |
| 2 | HUM-2 | 10.67 | 59.67 | 3.97 | 2.49 |
| 3 | MUM-2 | 8.6 | 62.47 | 6.11 | 2.94 |
| 4 | Paity Mung | 8.65 | 60.17 | 2.99 | 5.07 |
| 5 | Pant M-2 | 9.43 | 61.25 | 3.67 | 2.17 |
| 6 | Pant M-4 | 8.8 | 61.75 | 2.19 | 2.95 |
| 7 | PKV AKM-4 | 8.21 | 61.83 | 2.82 | 2.49 |
| 8 | Pusa-0672 | 9.17 | 61.31 | 2.98 | 2.95 |
| 9 | Pusa-9072 | 8.81 | 62.05 | 2.9 | 3.17 |
| 10 | RMG-62 | 8.53 | 63.56 | 3.97 | 2.48 |
| 11 | RMG-268 | 9.23 | 61.99 | 3.43 | 2.94 |
| 12 | Samrat | 7.83 | 62.09 | 4.47 | 2.98 |
| | SEm± | 0.11 | 0.1 | 0.02 | 0.02 |
| | CD at 5% | 0.33 | 0.3 | 0.06 | 0.06 |

Table: 2 Values of Reducing Sugar (%), Crude Protein (%), Soluble Protein (%), Methionine (g/16g N) in 12 Mung bean Cultivars

| S. N. | Cultivar/Variety | Reducing Sugar (%) | Crude Protein (%) | Soluble Protein (%) | Methionine (g/16g N) |
|-------|-------------------|--------------------|-------------------|---------------------|----------------------|
| 1 | BPMR-145 | 0.67 | 23.98 | 18.83 | 1 |
| 2 | HUM-2 | 0.92 | 25.16 | 15.48 | 1 |
| 3 | MUM-2 | 0.66 | 24.79 | 20.62 | 1.95 |
| 4 | Paity Mung | 0.61 | 24.81 | 16.8 | 0.79 |
| 5 | Pant M-2 | 0.8 | 24.51 | 17.26 | 1.22 |
| 6 | Pant M-4 | 0.66 | 23.95 | 18.07 | 0.91 |
| 7 | PKV AKM-4 | 0.64 | 23.8 | 18.34 | 0.99 |
| 8 | Pusa-0672 | 0.39 | 24.97 | 15.44 | 1.18 |

| | | | | | |
|----|------------------|------|-------|-------|------|
| 9 | Pusa-9072 | 0.61 | 24.81 | 17.33 | 1.14 |
| 10 | RMG-62 | 0.74 | 24.16 | 15.77 | 0.64 |
| 11 | RMG-268 | 0.77 | 24.31 | 18.84 | 0.94 |
| 12 | Samrat | 0.8 | 23 | 16.49 | 0.98 |
| | SEm± | 0 | 0.23 | 0.04 | 0.03 |
| | CD at 5% | 0.01 | 0.66 | 0.13 | 0.08 |

Table: 3 Values of Crude Fat Content (%), Calorific Value (kcal/100g), Total Phenol (mg/100g), Ash (%) in 12 Mung bean Cultivars

| S. N. | Cultivar/Variety | Crude Fat Content (%) | Calorific Value (kcal/100g) | Total Phenol (mg/100g) | Ash (%) |
|-------|-------------------|-----------------------|-----------------------------|------------------------|---------|
| 1 | BPMR-145 | 1.15 | 354.86 | 67.34 | 3.74 |
| 2 | HUM-2 | 1.16 | 351.17 | 89.33 | 3.68 |
| 3 | MUM-2 | 0.95 | 360.52 | 75.46 | 3.63 |
| 4 | Paity Mung | 1.22 | 351.16 | 72.63 | 3.5 |
| 5 | Pant M-2 | 1.03 | 355.26 | 71.77 | 4.05 |
| 6 | Pant M-4 | 1.02 | 351.59 | 73.98 | 3.41 |
| 7 | PKV AKM-4 | 1.12 | 358.15 | 62.31 | 3.62 |
| 8 | Pusa-0672 | 1.27 | 356.09 | 60.18 | 3.58 |
| 9 | Pusa-9072 | 1.51 | 351.59 | 74.83 | 3.95 |
| 10 | RMG-62 | 1.04 | 356.59 | 65.46 | 3.61 |
| 11 | RMG-268 | 1.01 | 356.08 | 71.33 | 3.45 |
| 12 | Samrat | 1.28 | 349.14 | 80.16 | 3.65 |
| | SEm± | 0.04 | 1 | 0.1 | 0.06 |
| | CD at 5% | 0.13 | 2.93 | 0.29 | 0.17 |

Table: 4 Values of Fe Content (mg/100g), Cu Content (mg/100g), Zn Content(mg/100g), in 12 Mung beanCultivars

| S. N. | Cultivar/Variety | Fe Content (mg/100g) | Cu Content(mg/100g) | Zn Content(mg/100g) |
|-------|-------------------|----------------------|---------------------|---------------------|
| 1 | BPMR-145 | 14.89 | 1.69 | 3.56 |
| 2 | HUM-2 | 11.49 | 2.12 | 3.25 |
| 3 | MUM-2 | 12.19 | 1.72 | 3.11 |
| 4 | Paity Mung | 13.81 | 1.71 | 3.32 |
| 5 | Pant M-2 | 11.39 | 1.87 | 3.27 |
| 6 | Pant M-4 | 13.71 | 1.9 | 3.38 |
| 7 | PKV AKM-4 | 14.12 | 2.05 | 3.4 |
| 8 | Pusa-0672 | 10.32 | 1.63 | 2.55 |
| 9 | Pusa-9072 | 14.28 | 3.28 | 3.49 |
| 10 | RMG-62 | 12.43 | 3 | 2.78 |
| 11 | RMG-268 | 11.27 | 1.64 | 3.43 |
| 12 | Samrat | 10.45 | 1.55 | 2.78 |
| | SEm± | 0.17 | 0.04 | 0.06 |
| | CD at 5% | 0.5 | 0.11 | 0.18 |

Conclusion

Therefore, it is possible to propose that mung bean seedlings are responsible for alterations in the nutritional value, biochemical makeup, growth parameters, and pigments involved in photosynthetic processes. In comparison to the dry mung bean seeds, there was a significant increase in proteins and amino acids due to the de novo synthesis of new proteins and the accumulation of some existing proteins. Enhancing the nutritional value of legumes through these methods is crucial for mitigating the food crisis. Therefore, increasing the intake of germinated mung beans is advised in order to improve nutrition absorption and prevent disease.

Reference

1. Sathyamoorthi K, Amanullah M M, Vaiyapuri K & Somasundaram E (2017) Res J Agric Biosci, 3(6), 886. Anonymous (2018) Package of Practices for Crops of Punjab, Punjab Agricultural University, Ludhiana. Khan M R I, Gatehouse J A & Boulter D (2019) J Exp Bot, 31, 1599.
2. Sekhon H S, Singh G, Brar J S, Shangmugasundaram S, Bains T S & Kooner B S (2004) Technology Package for Mungbean Cultivation in Punjab (India), DFID-AVRDC PAU mungbean project .

3. Sital J S & Narang A S (2009) Proc Nat Acad Sci (India), 64, 189.
4. Sital J S, Phutela S, Kaur A, Kapur J & Narang A S (1997) J Pl Sci Res, 13, 38. Schroeder H E (2016) J Sci Food Agric, 33, 623.
5. Habibullah, M. A., Abbas, M. and Shah, H. U. (2007). Proximate and mineral composition of mungbean, *Sarhad J. Agri.*, **23** (2): 463-466.
6. Bhatti, N., Gilani, A. H. and Ahmad, N. (2000). Nutritional value of greengram [*Vigna radiata* (L.) wilczek] as affected by cooking and supplementation. *Archivos Latinoamericanos de Nutrition*, **50** (4): 374-379.
7. Butt, M. S. and Batool, R. (2010). Nutritional and functional properties of some promising legumes protein isolates. *Pak. J. Nutrition*. **9**: 373-379.
8. Blessing, I. A. and Gregory, I. O. (2010). Effect of processing on the proximate composition of the dehulled and undehulled mungbean (*Vigna radiata* (L.) Wilczek) flours. *Pakistan Journal of Nutrition*, **9** (10): 1006-1016.
9. Mubarak, A. E. (2005). Nutritional composition and antinutritional factors of mungbean seeds (*Phaseolus aureus*) as affected by some home traditional processes, *Food Chem.* **89**: 489-495.
10. Tanusi, S., Kasai, T. and Kawamura, S. (1972). Determination of oligosaccharides in some edible legume seeds. *Journal of Japanese Society of Food Nutrition*. **25**: 25.
11. Chakraborty, R. (1993). Biochemical compositions of matured seeds of green gram [*Vigna radiata* (L.) Wilczek] in relation to cooking. M.Sc. (Agri.) Thesis, Assam Agricultural University, Jorhat. p. 34-77.
12. Kakati, P., Deka S. C., Kotoki, D. and Saikia, S. (2010). Effect of traditional methods of processing on the nutrient contents and some antinutritional factors in newly developed cultivars of green gram [*Vigna radiata* (L.) Wilczek] and black gram (*Vigna mungo* (L.) Hepper] of Assam, India. *International Food Research Journal*. **17**: 377-384.
13. Anonymous (2012). Report. Division of Basic Sciences. ICAR-Indian Institute of Pulses Research, Kanpur, India.
14. Mondal, N. K., Datta, J. K. and Banerjee, A. (2013). Biochemical response of mungbean (*Vigna radiata* L.) under the influence of reduced dose of chemical fertilizer and different time and method of application of biofertilizer. *Journal of Agricultural Technology*. **9** (3): 643-658.
15. Naivikul, O. and D'apponia. (1976). Comparison of legume and wheat flour carbohydrate. I. Sugar analysis. *American Association of Cereal Chemists*. **55** (6): 913-918.

16. Adel, A., Shehata, Y, and Thannoun, A. M. (1980). Chemical and amino acid composition of Iraqi mungbeans. *Zeitschrift fur Lebensmittel-Untersuchung und Forschung A*,**171** (5): 360-362.
17. Paul, T., Mozumder, N. H. M., Sayed, M. A. and Akhtaruzzaman, A. (2011). Proximate composition, mineral content and determination of protease activity from greengram (*Vigna radiata* L.Wilczek). *Bangladesh ResearchPublication Journal*. **5** (3): 207-213.